

Family Gauge Bosons with an Inverted Mass Hierarchy

Yoshio Koide^{a,b} and Toshifumi Yamashita^b

^a *Department of Physics, Osaka University, Toyonaka, Osaka 560-0043, Japan*

E-mail address: koide@het.phys.sci.osaka-u.ac.jp

^b *MISC, Kyoto Sangyo University, Kyoto 603-8555, Japan*

E-mail address: tyamashi@cc.kyoto-su.ac.jp

Abstract

A model that gives family gauge bosons with an inverted mass hierarchy is proposed, stimulated by Sumino's cancellation mechanism for the QED radiative correction to the charged lepton masses. The Sumino mechanism cannot straightforwardly be applied to SUSY models because of the non-renormalization theorem. In this paper, an alternative model which is applicable to a SUSY model is proposed. It is essential that family gauge boson masses $m(A_i^j)$ in this model is given by an inverted mass hierarchy $m(A_i^i) \propto 1/\sqrt{m_{ei}}$, in contrast to $m(A_i^i) \propto \sqrt{m_{ei}}$ in the original Sumino model. Phenomenological meaning of the model is also investigated. In particular, we notice a deviation from the e - μ universality in the tau decays.

PACS numbers: 11.30.Hv, 12.60.-i, 12.60.Jv

1 Introduction

It seems to be meaningful to consider that the flavor physics can be understood on the basis of a family symmetry [1]. Regrettably, since a constraint from the observed K^0 - \bar{K}^0 mixing is very tight, the family gauge boson masses must be super heavy, so that it is hard to observe such gauge boson effects in terrestrial experiments. Recently, one positive effect of the existence of family gauge bosons has been pointed out by Sumino [2, 3]. Since we propose a model with a Sumino-like mechanism in the present paper, we first give a brief review of the Sumino mechanism.

We know a miraculous formula for the charged lepton masses [4]:

$$K \equiv \frac{m_e + m_\mu + m_\tau}{(\sqrt{m_e} + \sqrt{m_\mu} + \sqrt{m_\tau})^2} = \frac{2}{3}, \quad (1.1)$$

which is satisfied with the order of 10^{-5} for the pole masses, i.e. $K^{pole} = (2/3) \times (0.999989 \pm 0.000014)$ [5]. However, in conventional flavor models, “masses” do not mean “pole masses”, but “running masses”. The formula (1.1) is only valid with the order of 10^{-3} for the running masses, e.g. $K(\mu) = (2/3) \times (1.00189 \pm 0.00002)$ at $\mu = m_Z$. The deviation of $K(\mu)$ from K^{pole} is caused by a logarithmic term $m_{ei} \log(\mu/m_{ei})$ in the QED radiative correction term [6]

$$m_{ei}(\mu) = m_{ei}^{pole} \left[1 - \frac{\alpha_{em}(\mu)}{\pi} \left(1 + \frac{3}{4} \log \frac{\mu^2}{m_{ei}^2(\mu)} \right) \right]. \quad (1.2)$$

Note that the value of K is invariant under the transformation

$$m_{ei} \rightarrow m_{ei}(1 + \varepsilon_0 + \varepsilon_i), \quad (1.3)$$

when $\varepsilon_i = 0$, where ε_0 and ε_i are factors which are independent of and dependent on the family-number i ($i = 1, 2, 3$), respectively. If the logarithmic term in the radiative correction (1.2) due to the photon can be cancelled by a some additional effect, the relation $K^{pole} = K(\Lambda) \equiv 2/3$ can be satisfied (Λ is an energy scale at which $K = 2/3$ is given).

Sumino [2] has seriously taken why the mass formula $K = 2/3$ is so remarkably satisfied with the pole masses, and proposed a cancellation mechanism of the logarithmic term in the radiative correction (1.2). He has assumed that the family symmetry is local, and that the logarithmic term is canceled by that due to the family gauge bosons. In the Sumino model, the left- and right-handed charged leptons e_{Li} and e_{Ri} are assigned to $\mathbf{3}$ and $\mathbf{3}^*$ of a U(3) family symmetry, respectively. (A similar fermion assignment has been proposed by Appelquist, Bai and Piai [7].) The charged lepton mass term is generated by a would-be Yukawa interaction

$$H_e = \frac{y_e}{\Lambda^2} \bar{\ell}_L^i \Phi_{i\alpha}^e \Phi_{\alpha j}^{eT} e_R^j H, \quad (1.4)$$

where i and α are indices of U(3) and O(3), respectively, H is the Higgs scalar in the standard non-SUSY model, and $\ell_L = (\nu, e^-)$. The VEV matrix $\langle \Phi_e \rangle$ is assumed as

$$\langle \Phi_e \rangle = \text{diag}(v_1, v_2, v_3) \propto \text{diag}(\sqrt{m_e}, \sqrt{m_\mu}, \sqrt{m_\tau}). \quad (1.5)$$

Then, the family gauge boson masses $m(A_i^j)$ are given by

$$m^2(A_j^i) \propto \langle \Phi_{i\alpha}^e \rangle \langle (\Phi^e)^{\dagger\alpha i} \rangle + \langle \Phi_{j\alpha}^e \rangle \langle (\Phi^e)^{\dagger\alpha j} \rangle \propto m_{ei} + m_{ej}. \quad (1.6)$$

Since in the Sumino model, the charged lepton fields (e_L, e_R) are assigned to $(e_L, e_R) \sim (\mathbf{3}, \mathbf{3}^*)$ of the U(3), the family gauge bosons in the off-diagonal elements A_i^j ($i \neq j$) cannot contribute to the radiative corrections. Then, the cancellation takes place between $\log m_{ei}$ in the QED diagram and $\log m(A_i^i) \propto \log m_{ei}$ in the family gauge boson diagram. (Of course, the family gauge boson coupling constant g_F must satisfy a relation $g_F^2/4 \simeq e^2$.) As a result, we can obtain $K(\Lambda) = K^{pole}$.

In Ref.[2], and also in this paper, it is assumed that the formula (1.1) is already given at an energy scale Λ , and is not discussed how to derive the formula¹.

Now, it is interesting to apply this Sumino mechanism to a supersymmetric (SUSY) scenario. It should be noted that a vertex correction is, in general, vanishing in the SUSY scenario, so that the Sumino mechanism cannot be applied to SUSY models straightforwardly. In this paper, we investigate how to restore the Sumino mechanism in such models. The essential idea in the present model is as follows: The cancellation in the Sumino mechanism occurs due to the vertex correction diagram, while that in the present paper does due to the wave function renormalization diagrams. (The details are given in Sec.2.) In the original Sumino mechanism,

¹The first attempt to understand the mass formula (1.1) from the bilinear form of Φ_e has been done by assuming a U(3) family symmetry and a ‘‘nonet’’ ansatz for Φ [8]. For more plausible derivation of the formula, see Ref. [3], where the model is based on a U(9) family symmetry.

the QED correction is cancelled by $g_F^L g_F^R \log m(A_i^i) \propto -g_F^2 \log m_{ei}$ with $g_F^L = -g_F^R \equiv g_F$, while in the present SUSY model it is done by $(g_F^{L,R})^2 \log m(A_i^j)$. For this purpose, we will assume the gauge boson masses with an inverted hierarchy $m^2(A_i^i) \propto 1/m_{ei}$ as we state in the next section.

In the Sumino model, the gauge boson A_j^i couples to

$$(J_\mu^{Sumino})_i^j = \bar{\psi}_L^j \gamma_\mu \psi_{Li} - \bar{\psi}_{Ri} \gamma_\mu \psi_R^j, \quad (1.7)$$

so that the current-current interactions inevitably cause interactions which violate the individual family number N_F by $|\Delta N_F| = 2$. This $|\Delta N_F| = 2$ effects are somewhat troublesome in the low energy phenomenology. In contrast to the Sumino currents (1.7), our family currents are given by the canonical form

$$(J_\mu)_i^j = \bar{\psi}_L^j \gamma_\mu \psi_{Li} + \bar{\psi}_R^j \gamma_\mu \psi_{Ri} = \bar{\psi}^j \gamma_\mu \psi_i, \quad (1.8)$$

so that the $|\Delta N_F| = 2$ effects appear only through a small quark family mixing.

In summary, the present model has the following characteristics compared with the Sumino model: (i) Since we can assign the family multiplets as $(f_L, f_R) \sim (\mathbf{3}, \mathbf{3})$ of the U(3), it is easy to make the model anomaly-less. (ii) In contrast to the Sumino currents (1.7), we can take a canonical form of the family currents (1.8). Therefore, the $|\Delta N_F| = 2$ effects appear only through a small quark mixing. (iii) Family gauge bosons with the lowest and highest masses are A_3^3 and A_1^1 , respectively. Note that gauge bosons which can couple to the light quarks u , d and s are only A_j^i with $i, j = 1, 2$. Therefore, we may consider that the contributions of the family gauge boson exchanges with $i, j = 1, 2$ are reduced compared with a conventional family gauge boson model. This means that we can take a lower value of $m(A_3^3)$ (e.g. a few TeV), and may expect to observe $A_3^3 \rightarrow \tau^+ \tau^- / b\bar{b} / t\bar{t}$ via $b\bar{b} / t\bar{t}$ associated production in the LHC. In Sec.4, we will investigate a deviation from the e - μ universality in the tau lepton decays. The present data allow the lowest gauge boson mass $m(A_3^3)$ to be of the order of a few TeV. We will also investigate possible family-number conserving but lepton flavor violating decays of K , D and B mesons. The observations of $K \rightarrow \pi \mu^+ e^-$ and $B \rightarrow K \mu^- \tau^+$ are within our reach. Thus, we can expect fruitful low energy phenomenology.

2 Cancellation mechanism in a SUSY model

In a SUSY model, the contributions of the family gauge bosons in the vertex correction diagram become vanishing, so that the original Sumino mechanism does not work. On the other hand, those from the wave function renormalization diagram still remain:

$$\delta m_{ei} = 2m_{ei} \sum_j \frac{\gamma_{eij}}{(4\pi)^2} \log \frac{\mu}{M_{ij}} = m_{ei} \frac{\alpha_F}{2\pi} \sum_j \log \frac{M_{ij}^2}{\mu^2}. \quad (2.1)$$

Here γ_{eij} gives the anomalous dimension γ_{ei} when summed over j ,

$$\gamma_{eij} = -2g_F^2 \sum_a (T^a)_{ij} (T^a)_{ji}, \quad \gamma_{ei} = \sum_j \gamma_{eij}, \quad (2.2)$$

where T^a is the generator of the U(3). Therefore, in the present model, the values of ε_i defined

by Eq.(1.3) are given by

$$\varepsilon_i = \rho \left(\log \frac{m_{ei}^2}{\mu^2} + \zeta \sum_j \log \frac{M_{ij}^2}{\mu^2} \right), \quad \rho = \frac{3}{4} \frac{\alpha_{em}}{\pi}, \quad \zeta = \frac{2}{3} \frac{\alpha_F}{\alpha_{em}}, \quad (2.3)$$

and M_{ij} are the family gauge boson masses $M_{ij} = m(A_i^j)$ given by

$$M_{ij}^2 \propto \frac{1}{m_{ei}} + \frac{1}{m_{ej}}. \quad (2.4)$$

(For a model for M_{ij} , see the next section.)

Note that since the gauge bosons A_i^j with $j \neq i$ can contribute to the ε_i term, differently from the Sumino model, the QED $\log m_{ei}$ term cannot be canceled by the gauge boson terms exactly even if we adjust the parameter ζ . The ratio of $K(m_{ei})$ to $K(m_{ei}^0)$ for $m_{ei} = m_{ei}^0(1 + \varepsilon_0 + \varepsilon_i)$

$$R \equiv \frac{K(m_{ei})}{K(m_{ei}^0)}, \quad (2.5)$$

is, in general, dependent on the values ζ and ε_0 . (The value of ε_0 is practically not essential as $|\varepsilon_0| \ll 1$.)

Next, we investigate the ζ -dependence of the ratio R . Since the ε_0 term can always be shifted by a common value, we shift the ε_i terms as $\varepsilon_i \rightarrow \varepsilon_i - \varepsilon_3$ and $\varepsilon_0 \rightarrow \varepsilon_0 + \varepsilon_3$. Then, we obtain

$$\frac{1}{\rho} \varepsilon_1 = \log \frac{m_{e1}^2}{m_{e3}^2} + \zeta \left(\frac{1}{2} \log \frac{m_{e3}^2}{m_{e1}^2} + \frac{1}{2} \log \frac{m_{e2}^2}{m_{e1}^2} + \log \frac{1 + m_{e1}/m_{e2}}{1 + m_{e2}/m_{e3}} \right), \quad (2.6)$$

$$\frac{1}{\rho} \varepsilon_2 = \log \frac{m_{e2}^2}{m_{e3}^2} + \zeta \left(\frac{1}{2} \log \frac{m_{e3}^2}{m_{e2}^2} + \log \frac{1 + m_{e1}/m_{e2}}{1 + m_{e1}/m_{e3}} \right), \quad (2.7)$$

and $\varepsilon_3/\rho = 0$. Here, the first terms in the parentheses in the right hand sides represent the contributions of the diagonal gauge fields A_i^i , while the succeeding terms do those of the off-diagonal ones A_i^j ($j \neq i$). As expected, by setting ζ appropriately ($\zeta = 2$), the diagonal gauge fields cancel the QED logarithmic terms as in the Sumino mechanism, but the off-diagonal ones make the cancellation incomplete, as $\varepsilon_2/\rho = 2 \log[(1 + m_{e1}/m_{e2})/(1 + m_{e1}/m_{e3})] \simeq 2(m_{e1}/m_{e2})$ and $\varepsilon_1/\rho \simeq \log(m_{e1}/m_{e2})^2$ in this case. Interestingly, however, since ε_2 is quite small and safely neglected, the effect on the parameter K which has a mild dependence on m_{e1} is relatively suppressed. Although the suppression is not sufficient, $R - 1 = \mathcal{O}(10^{-4})$, we expect that the deviation is cancelled by some other effects, such as the tau-Yukawa effect, a misarrangement of ζ for instance due to the renormalization group (RG) effects, and so on.

If we want more precise value of ζ at which R becomes 1, we can obtain it numerically. For convenience, we use the following input values: the observed charged lepton pole masses [5]

$m_{e1}^0 = 0.510998910 \times 10^{-3}$ GeV, $m_{e2}^0 = 0.105658367$ GeV, $m_{e3}^0 = 1.77682$ GeV; the fine structure constant at $\mu = m_Z$, $\alpha(m_Z) = 1/127.916$ [5]. For the time being, we do not specify an energy scale Λ at which the formula $K(\Lambda) = 2/3$ holds. Then, we find that a value of ζ which gives $R = 1$ is $\zeta = 1.752$. We can check that the value of $R - 1$ at $\zeta = 1.752$ is insensitive of the value ε_0 . The value $\zeta = 1.752$ is near to a value $7/4$. If we consider $\zeta = 7/4$, we can also show that $R - 1$ is always smaller than 10^{-5} independently of the value ε_0 . Thus, although the present model cannot give rigorous cancellation of the $\log m_{ei}$ term, it can practically give $R = 1$ with an accuracy of 10^{-5} .

3 Model

A simple way to make a model for the charged leptons anomaly-less is to assign the lepton doublet ℓ and charged lepton singlet e^c to $\mathbf{3}$ and $\mathbf{3}^*$ of the U(3) family symmetry, respectively, in contrast to the Sumino model [2, 3] where both have been assigned to $\mathbf{3}$. If we adopt a yukawaon model[9], the would-be Yukawa interaction for the charged lepton sector is given by $(y_e/\Lambda)\ell_i(Y_e)_j^i e^{cj} H_d$. Then, however, another term $(y'_e/\Lambda)\ell_i(Y_e)_j^j e^{ci} H_d$ is also allowed by the symmetry, and there are no reasons to forbid it. (This problem always appear when we take a model with $\ell \sim \mathbf{3}$ and $e^c \sim \mathbf{3}^*$.) Therefore, in the present paper, we do not adopt such a yukawaon model. Following to the Sumino model, we will consider a bilinear contribution $(\bar{\Phi}_e)_\alpha^i (\Phi_e)_j^\alpha$ instead of $(Y_e)_j^i$. To be more concrete, we assign $(\bar{\Phi}_e)_\alpha^i \sim (\mathbf{3}^*, \mathbf{3})$ and $(\Phi_e)_j^\alpha \sim (\mathbf{3}, \mathbf{3}^*)$ of U(3) \times U(3)' family symmetries. Note that we consider U(3)' instead of O(3) in the Sumino model. This allows us to take a flavor basis in which $\langle \Phi_e \rangle$ is diagonal. Hereafter, we simply denote Φ_e and $\bar{\Phi}_e$ in the present model as Φ and $\bar{\Phi}$, respectively.

We take the following would-be Yukawa interaction terms

$$W_Y = y_\ell \ell_i \bar{\Phi}_\alpha^i \bar{L}^\alpha + y_{Hd} L_\alpha H_d E^{c\alpha} + y_e \bar{E}_\alpha^c \Phi_j^\alpha e^{cj} + M_E E^{c\alpha} \bar{E}_\alpha^c + M_L L_\alpha \bar{L}^\alpha, \quad (3.1)$$

where $\ell_i = (\nu_i, e_i)$, e^{ci} , $L_\alpha = (N_\alpha, E_\alpha)$, $E^{c\alpha}$, $\bar{L}^\alpha = (\bar{N}^\alpha, \bar{E}^\alpha)$ and \bar{E}_α^c , have the electric charges $(0, -1)$, $+1$, $(0, -1)$, $+1$, $(0, +1)$ and -1 , respectively. Here, (E^c, \bar{E}^c) and (L, \bar{L}) are vector-like SU(2) $_L$ singlets and doublets, respectively. Then, we obtain the following effective superpotential

$$W_Y^{eff} = \frac{y_{Hd} y_\ell y_e}{M_E M_L} \ell_i \bar{\Phi}_\alpha^i \Phi_j^\alpha e^{cj} H_d. \quad (3.2)$$

Note that the counterpart of the y' -term $\text{Tr}[\bar{\Phi}\Phi]\ell_i e^{ci} H_d$ is not generated at the tree level, and then, in great contrast to non-SUSY models, is protected against the radiative corrections thanks to the nonrenormalization theorem. To be complete, we should forbid a term $\ell_i e^{ci} H_d$ and hopefully the above effective nonrenormalizable term generated already at the cutoff scale Λ . The former term can be forbidden by a U(1) $_S$ symmetry as usual. A concrete example of assignment of the charge and other quantum numbers (including those of the U(1) $_R$ symmetry that the superpotential considered here has) are shown in Table 1. The latter term may be forbidden effectively by assuming that the cutoff scale is large enough, or by replacing for instance the mass M_E with a VEV of a field S charged under the U(1) $_S$ symmetry. Here, we employ the former way for simplicity and consider only renormalizable terms, while we also introduce the field S for the later convenience.

To avoid the massless Nambu-Goldstone mode, we assume that the $U(1)_S$ charge conservation is broken by a soft term

$$W_{br} = \mu_S S \theta_S - \varepsilon \mu_S^2 \theta_S, \quad (3.3)$$

where S and θ_S are family singlets, and ε has been put in order to denote that the R charge conservation in the term (3.3) is softly broken with a small extent ε . The superpotential (3.3) leads to $\langle S \rangle = \varepsilon \mu_S$.

The effective superpotential (3.2) reduces to the charged lepton Yukawa interaction when Φ and $\bar{\Phi}$ acquire VEVs. The magnitude of $\langle \Phi \rangle \langle \bar{\Phi} \rangle$ is set by

$$W_\Phi = \lambda_1 \Phi_i^\alpha \bar{\Phi}_\alpha^i \theta_\Phi - \lambda_2 S^2 \theta_\Phi, \quad (3.4)$$

where θ_Φ is a family singlet field, which leads to

$$\text{Tr}[\langle \Phi \rangle \langle \bar{\Phi} \rangle] = \varepsilon^2 \frac{\lambda_2}{\lambda_1} \mu_S^2. \quad (3.5)$$

As mentioned, we do not discuss how appropriate forms (1.5) of the VEVs, $\langle \Phi \rangle$ and $\langle \bar{\Phi} \rangle$, are obtained but just assume that a superpotential $W_K(\Phi, \bar{\Phi}, \dots)$ leads them. Here, in addition, we assume that W_K is invariant² under the exchange of the $U(3)$ and the $U(3)'$, which exchanges Φ and $\bar{\Phi}$, and that the VEVs respect this $S_2 (= Z_2)$ symmetry:

$$\langle \Phi \rangle = \langle \bar{\Phi} \rangle = v_\Phi Z = v_\Phi \text{diag}(z_1, z_2, z_3). \quad (3.6)$$

Here, in the last equality, we have used the $U(3)$ and $U(3)'$ degrees of freedom to diagonalize $\langle \Phi \rangle$, and the parameters z_i are normalized as $z_1^2 + z_2^2 + z_3^2 = 1$, without losing generality. With this notation, from Eq.(3.5), we obtain

$$v_\Phi^2 = \varepsilon^2 \frac{\lambda_2}{\lambda_1} \mu_S^2. \quad (3.7)$$

Next, we investigate a superpotential for the field Ψ whose VEV $\langle \Psi \rangle$ gives an inverted mass hierarchy (2.4):

$$\begin{aligned} W_{\Phi\Psi} = & (\lambda_A \bar{\Psi}_\alpha^i \Phi_j^\alpha + \bar{\lambda}_A \bar{\Phi}_\alpha^i \Psi_j^\alpha) (\Theta_A)_i^j + (\lambda'_A \bar{\Psi}_\alpha^i \Phi_i^\alpha + \bar{\lambda}'_A \bar{\Phi}_\alpha^i \Psi_i^\alpha - \mu_A S) (\Theta_A)_j^i \\ & + (\lambda_B \Phi_i^\alpha \bar{\Psi}_\beta^i + \bar{\lambda}_B \Psi_i^\alpha \bar{\Phi}_\beta^i) (\Theta_B)_\alpha^\beta + (\lambda'_B \Phi_i^\alpha \bar{\Psi}_\alpha^i + \bar{\lambda}'_B \Psi_i^\alpha \bar{\Phi}_\alpha^i - \mu_B S) (\Theta_B)_\beta^i. \end{aligned} \quad (3.8)$$

Again, we impose that $W_{\Phi\Psi}$ is S_2 invariant, i.e. $\lambda_A = \bar{\lambda}_B$, $\bar{\lambda}_A = \lambda_B$, $\lambda'_A = \bar{\lambda}'_B$, $\bar{\lambda}'_A = \lambda'_B$ and $\mu_A = \mu_B$. Then, from the F -flatness conditions, we obtain

$$\langle \Psi \rangle = \langle \bar{\Psi} \rangle = v_\Psi Z^{-1}, \quad (3.9)$$

which satisfies the D -term condition too. We also obtain

$$v_\Phi v_\Psi = 3\varepsilon \frac{\mu_S \mu_A}{\lambda_A + 3\lambda'_A + \bar{\lambda}_A + 3\bar{\lambda}'_A}. \quad (3.10)$$

²The superpotential (3.1) does not possess this invariance.

	ℓ	e^c	H_d	L	\bar{L}	E^c	\bar{E}^c	Φ	$\bar{\Phi}$	Ψ	$\bar{\Psi}$	θ_Φ	Θ_A	Θ_B	S	θ_S
$SU(2)_L$	2	1	2	2	2	1	1	1	1	1	1	1	1	1	1	1
$U(3)$	3	3*	1	1	1	1	1	3	3*	3	3*	1	8+1	1	1	1
$U(3)'$	1	1	1	3	3*	3*	3	3*	3	3*	3	1	1	8+1	1	1
$U(1)_S$	0	0	-1	1	-1	0	-1	1	1	0	0	-2	-1	-1	1	-1
$U(1)_R$	1	1	0	1	1	1	1	0	0	0	0	2	2	2	0	2

Table 1: The fields in the present model and their quantum numbers.

Comparing with Eq.(3.7), we see

$$\frac{v_\Phi}{v_\Psi} = \varepsilon k \sim O(\varepsilon), \quad (3.11)$$

where $k = (1/3)(\lambda_2/\lambda_1)(\lambda_A + 3\lambda'_A + \bar{\lambda}_A + 3\bar{\lambda}'_A)(\mu_S/\mu_A)$. Since the charged lepton masses m_{ei} are given by

$$M_e \equiv \text{diag}(m_{e1}, m_{e2}, m_{e3}) \propto v_\Phi^2 Z^2 = v_\Phi^2 \text{diag}(z_1^2, z_2^2, z_3^2), \quad (3.12)$$

from Eq.(3.2), the parameter values of z_i are given by

$$z_i = \frac{\sqrt{m_{ei}}}{\sqrt{m_{e1} + m_{e2} + m_{e3}}}, \quad (3.13)$$

where $(m_{e1}, m_{e2}, m_{e3}) = (m_e, m_\mu, m_\tau)$. The explicit values of z_i are given by

$$(z_1, z_2, z_3) = (0.016473, 0.23688, 0.97140). \quad (3.14)$$

Thus, we can approximately estimate the family gauge boson masses $m(A_j^i)$ as follows

$$\begin{aligned} M_{ij}^2 \equiv m^2(A_j^i) &= \frac{1}{2} g_F^2 \left[\sum_\alpha (\langle (\Psi^\dagger)_\alpha^i \rangle \langle \Psi_i^\alpha \rangle + \langle \bar{\Psi}_\alpha^i \rangle \langle (\bar{\Psi}^\dagger)_i^\alpha \rangle) + (i \rightarrow j) + O(\varepsilon^2) \right] \\ &\simeq g_F^2 v_\Psi^2 \left(\frac{1}{z_i^2} + \frac{1}{z_j^2} \right) \propto \left(\frac{1}{m_{ei}} + \frac{1}{m_{ej}} \right), \end{aligned} \quad (3.15)$$

if the mixing between the $U(3)$ gauge boson and the $U(3)'$ one can be neglected. This happens when the latter gauge boson is sufficiently heavy and we assume such a case. Namely, we assume another sector that breaks $U(3)'$ at a high scale which is basically decoupled from the sector we have discussed above.

In this case, in addition to the interactions in the superpotential (3.1), the gauge interaction (below the $U(3)'$ breaking scale) violates the S_2 symmetry assumed in W_K and $W_{\Phi\Psi}$, and the RG effects modify the S_2 relations shown above Eq.(3.9). Nevertheless, amazingly, the nonrenormalization theorem protects the VEV relations (3.6) and (3.9) against the RG effects, which justifies the above discussion.

So far, we have not discussed the neutrino mass matrix, because the purpose of the present paper is to discuss how to apply the Sumino mechanism to a SUSY model. Here, we would like to give a brief comment on the neutrino mass matrix. In order to obtain neutrino masses, we add, for instance, a new field N^c , which is $SU(2)_L$ doublet with $\mathbf{3}^*$ of the family symmetry $SU(3)'$. (However, we do not consider the vector-like partner \bar{N}^c unlike the case (E^c, \bar{E}^c) . Therefore, in order to make the model anomaly free, some additional fields are needed. In this paper, we do not comment on it.) The field generates the following superpotential terms in addition to Eq.(3.1):

$$y_{Hu}L_\alpha H_u N^{c\alpha} + y_M N^{c\alpha} (Y_M)_{\alpha\beta} N^{c\beta}. \quad (3.16)$$

The superpotential terms (3.1) and (3.16) leads to the effective neutrino mass matrix as Eq.(3.2):

$$W_\nu^{eff} = \frac{y_{Hu}y_\ell}{y_M M_L^2} (\ell_i H_u) \bar{\Phi}_\alpha^i (\langle Y_M \rangle^{-1})^{\alpha\beta} \bar{\Phi}_\beta^j (\ell_j H_u). \quad (3.17)$$

Here, we consider that the VEV value $\langle Y_M \rangle_{\alpha\beta}$ breaks $SU(3)'$ symmetry at a higher scale Λ' ($\Lambda' \gg \Lambda$). This realizes the above assumption used in Eq.(3.15), and is a reason that we will not discuss $SU(3)'$ family symmetry gauge boson effects in the following low energy scale phenomenology.

4 Possible effects of the family gauge bosons

Since the gauge boson masses are given by Eq.(3.15), we obtain the following hierarchical structure:

$$\begin{aligned} 2g_F^2 v_\Psi^2 \frac{1}{z_1^2} &= M_{11}^2 \simeq 2M_{12}^2 \simeq 2M_{13}^2, \\ 2g_F^2 v_\Psi^2 \frac{1}{z_2^2} &= M_{22}^2 \simeq 2M_{23}^2, \\ 2g_F^2 v_\Psi^2 \frac{1}{z_3^2} &= M_{33}^2. \end{aligned} \quad (4.1)$$

The family currents in the Sumino model are given by Eq.(1.7), while, in the present model, those are given by

$$(J_\mu)_j^i = \bar{e}_L^i \gamma_\mu e_{Lj} + \bar{e}_R^i \gamma_\mu e_{Rj} + \dots = \bar{e}^i \gamma_\mu e_j + \dots, \quad (4.2)$$

where, for convenience, we have denoted only charged lepton sector explicitly. Note that the effective current-current interactions in the Sumino model induce $\Delta N_F = 2$ interactions, while those in the present model do not induce such $\Delta N_F = 2$ interactions. In the present model, however, since the family number is defined in a diagonal basis of the charged lepton mass matrix, in general, quark mixings appear, so that family number violating modes will be observed through $U_u \neq \mathbf{1}$ and $U_d \neq \mathbf{1}$, where U_q ($q = u, d$) is defined by $U_{Lq}^\dagger M_q U_{Rq} = D_q$ (D_q is a diagonal matrix). For example, family currents in the down-quark sector are given by

$$(J_\mu^{(d)})_j^i = (\bar{d}_L^0 \gamma_\mu d_{Lj}^0) + (\bar{d}_R^0 \gamma_\mu d_{Rj}^0) = (U_{Ld}^\dagger)_k^i (U_{Ld})_j^l (\bar{d}_L^k \gamma_\mu d_{Ll}) + (L \rightarrow R), \quad (4.3)$$

so that the effective Hamiltonians for semileptonic modes and nonleptonic modes, H_{SL} and H_{NL} , are given by

$$H_{SL}^{eff} = \sum_{i,j,k,l} \frac{G_{ij}}{\sqrt{2}} (U_d^\dagger)_k^i (U_d)_j^l (\bar{d}^k \gamma_\mu d_l) (\bar{e}^j \gamma^\mu e_i), \quad (4.4)$$

$$H_{NL}^{eff} = \sum_{i,j,k,l,m,n} \frac{G_{ij}}{\sqrt{2}} (U_d^\dagger)_k^i (U_d)_j^l (U_d^\dagger)_m^j (U_d)_i^n (\bar{d}^k \gamma_\mu d_l) (\bar{d}^m \gamma^\mu d_n), \quad (4.5)$$

respectively, where $G_{ij}/\sqrt{2} = g_F^2/2M_{ij}^2 \simeq z_j^2/2v_\Psi^2$, and, for simplicity, we have put $U_{Ld} = U_{Rd} \equiv U_d$. In this section, we investigate possible phenomenology of the flavor violating modes, and discuss the scale of the gauge bosons.

Usually, the most strict constraint comes from the observed K^0 - \bar{K}^0 mixing. However, this $\Delta N_F = 2$ transition occurs only via the down-quark mixing $U_d \neq \mathbf{1}$, so that the constraint is highly dependent on the quark mass matrix model. In this paper, we do not discuss a model about the quark mixing. Only modes that are independent of the quark mixing structures are pure leptonic decays $e_i \rightarrow e_j + \bar{\nu}_j + \nu_i$. Therefore, first, let us investigate these pure leptonic decays based on the present model. The effective interactions via the family gauge boson exchanges are given by

$$\frac{G_{ij}}{\sqrt{2}} (\bar{\nu}_i \gamma_\mu \nu_j) (\bar{e}_j \gamma^\mu e_i), \quad (4.6)$$

against the conventional weak interactions

$$\frac{G_F}{\sqrt{2}} (\bar{e}_j \gamma_\mu (1 - \gamma_5) \nu_j) (\bar{\nu}_i \gamma^\mu (1 - \gamma_5) e_i), \quad (4.7)$$

where $G_F/\sqrt{2} = g_W^2/8M_W^2 = 1/2v_W^2$ ($v_W = 246$ GeV). By using the Fierz transformation, we obtain effective coupling constants (for the definitions, see [10]) in the current-current interactions

$$g_{LL}^V = 1 + \epsilon_j, \quad g_{RR}^V = 0, \quad g_{LR}^S = -2\epsilon_j, \quad g_{RL}^S = 0, \quad (4.8)$$

where $\epsilon_j \simeq (1/4)z_j^2(v_W/v_\Psi)^2$, and we have considered a case that the observed neutrinos are Majorana types. The result (4.8) gives the decay parameters [10] $\eta = 0$, $\rho = 3/4$, $\delta = 3/4$ and $\xi \simeq 1 - 2\epsilon_j^2$. Regrettably, the results for η , ρ and δ are identical with those in the standard model (SM) and the deviation of ξ from $\xi^{SM} = 1$ is too small to observe. On the other hand, in relation to the branching ratios, we predict

$$R_\tau \equiv \frac{1 + \epsilon_\mu}{1 + \epsilon_e} = \left[\frac{B(\tau^- \rightarrow \mu^- \bar{\nu}_\mu \nu_\tau) f(m_e/m_\tau)}{B(\tau^- \rightarrow e^- \bar{\nu}_e \nu_\tau) f(m_\mu/m_\tau)} \right]^{1/2}, \quad (4.9)$$

where $f(x)$ has been defined by $f(x) = 1 - 8x^2 + 8x^6 - x^8 - 12x^4 \ln x^2$ and $f(m_e/m_\tau)/f(m_\mu/m_\tau) = 1.028215$. Since $\epsilon_e \simeq z_1^2 r^2/4 = 6.8 \times 10^{-5} r^2$ and $\epsilon_\mu \simeq z_2^2 r^2/4 = 1.4 \times 10^{-2} r^2$ [$r \equiv v_W/v_\Psi$], we expect a deviation $\Delta R_\tau \equiv R_\tau - 1 \simeq \epsilon_\mu$. Present experimental values [5] $B(\tau^- \rightarrow \mu^- \bar{\nu}_\mu \nu_\tau) = (17.39 \pm 0.04)\%$ and $B(\tau^- \rightarrow e^- \bar{\nu}_e \nu_\tau) = (17.82 \pm 0.04)\%$ give

$$R_\tau^{exp} = 1.0017 \pm 0.0016, \quad (4.10)$$

i.e. $\epsilon_\mu \simeq 0.0017 \pm 0.0016$. This result seems to be in favor of the inverted gauge boson mass hierarchy although it is just at 1σ level. (If the gauge boson masses take a normal hierarchy, R_τ

will show $R_\tau < 1$.) However, if we take the central value $\Delta R_\tau \sim 0.0017$, it means $r \sim 0.35$. This value corresponds to $v_\Psi \sim 0.7$ TeV which is ruled out by Kaon rare decays as we will see next. At present, we should not take the value (4.10) rigidly. If we speculate $r \sim 10^{-1}$ (v_Ψ of a few TeV), we may expect a sizable deviation $\Delta R_\tau \simeq \epsilon_\mu \simeq z_2^2 r^2 / 4 \sim 10^{-4}$ from the e - μ universality in the tau lepton decays. We expect that the observation $\Delta R_\tau \simeq 10^{-4}$ will be accomplished by a tau-factory in the near future.

Next, we direct our attention to family number conserved modes in the limit of no quark mixing. Predicted values for those modes are insensitive to the explicit values of U_d and U_u as far as they are not so large. In particular, we investigate rare decays of pseudo-scalar mesons $0^- \rightarrow 0^- + e_i + \bar{e}_j$ ($i \neq j$) with $\Delta N_F = 0$, e.g. $K^+ \rightarrow \pi^+ \mu^+ e^-$, $B^+ \rightarrow K^+ \tau^+ \mu^-$, and so on. (Since our currents are pure vectors, they cannot contribute to decays $0^- \rightarrow e_i + \bar{e}_j$.) When we neglect the CP violation effects and the electromagnetic mass difference of pseudo-scalar mesons, we can predict the following branching ratios:

$$B(K^+ \rightarrow \pi^+ \mu^+ e^-) \simeq 2z_1^4 r^4 \frac{1}{2|V_{us}|^2} B(K^+ \rightarrow \pi^0 \mu^+ \nu_\mu) = 4.88 \times 10^{-8} r^4, \quad (4.11)$$

$$B(K^0 \rightarrow \pi^0 \mu^+ e^-) \simeq \frac{1}{2} z_1^4 r^4 \frac{1}{2|V_{us}|^2} B(K^0 \rightarrow \pi^- \mu^+ \nu_\mu) = 9.82 \times 10^{-8} r^4, \quad (4.12)$$

$$B(D^+ \rightarrow \pi^+ \mu^- e^+) \simeq z_1^4 r^4 \frac{1}{2|V_{cs}|^2} B(D^+ \rightarrow \bar{K}^0 \mu^+ \nu_\mu) \frac{f(m_\pi/m_D)}{f(m_K/m_D)} = 5.83 \times 10^{-9} r^4, \quad (4.13)$$

$$B(D^0 \rightarrow \pi^0 \mu^- e^+) \simeq \frac{1}{2} z_1^4 r^4 \frac{1}{2|V_{cs}|^2} B(D^0 \rightarrow K^- \mu^+ \nu_\mu) \frac{f(m_\pi/m_D)}{f(m_K/m_D)} = 1.03 \times 10^{-9} r^4, \quad (4.14)$$

$$B(B^+ \rightarrow K^+ \mu^- \tau^+) \simeq z_2^4 r^4 \frac{1}{2|V_{cb}|^2} B(B^+ \rightarrow \bar{D}^0 \tau^+ \nu_\tau) \frac{f(m_K/m_B)}{f(m_D/m_B)} = 1.51 \times 10^{-2} r^4, \quad (4.15)$$

$$B(B^0 \rightarrow K^0 \mu^- \tau^+) \simeq z_2^4 r^4 \frac{1}{2|V_{cb}|^2} B(B^0 \rightarrow D^- \tau^+ \nu_\tau) \frac{f(m_K/m_B)}{f(m_D/m_B)} = 2.37 \times 10^{-2} r^4, \quad (4.16)$$

where r and $f(x)$ have been defined below Eq.(4.9). (For simplicity, we have used approximate relation in the limit of massless charged leptons. Therefore, the numerical results should not be taken rigidly.) In Eq.(4.12), under the approximation of neglecting CP violation, we read $B(K^0 \rightarrow \pi^- \mu^+ \nu_\mu)$ as $B(K_L \rightarrow \pi^\pm \mu^\mp \nu_\mu) = (1/2)B(K^0 \rightarrow \pi^- \mu^+ \nu_\mu) + (1/2)B(\bar{K}^0 \rightarrow \pi^+ \mu^- \bar{\nu}_\mu) = B(K^0 \rightarrow \pi^- \mu^+ \nu_\mu) = B(\bar{K}^0 \rightarrow \pi^+ \mu^- \bar{\nu}_\mu)$ (and also $B(K^0 \rightarrow \pi^0 \mu^+ e^-)$ as $B(K_L \rightarrow \pi^0 \mu^\pm e^\mp)$). In the numerical results in Eqs.(4.11) - (4.16), we have used the observed values [5] $B(K^+ \rightarrow \pi^0 \mu^+ \nu_\mu) = 3.353 \times 10^{-2}$, $B(K_L \rightarrow \pi^\mp \mu^\pm \nu_\mu) = 0.2704$, $B(D^+ \rightarrow \bar{K}^0 \mu^+ \nu_\mu) = 9.4 \times 10^{-2}$, $B(D^0 \rightarrow K^- \mu^+ \nu_\mu) = 3.31 \times 10^{-2}$, $B(B^+ \rightarrow \bar{D}^0 \tau^+ \nu_\tau) = 7 \times 10^{-3}$ and $B(B^0 \rightarrow D^- \tau^+ \nu_\tau) = 1.1 \times 10^{-2}$. For reference, we list the predicted values for $v_W/v_\Psi \sim 10^{-1}$ (and present experimental upper




Fig. 1 Predicted branching ratios $B(P \rightarrow P' e_i^+ e_j^-)$ versus the VEV value $v_F \equiv v_\Psi$. The marks \bullet and the dashed lines denote present lower limits of the observed branching ratios.

limits [5]) as follows:

$$\begin{aligned}
B(K^+ \rightarrow \pi^+ \mu^+ e^-) &\sim 5 \times 10^{-12} &< 1.3 \times 10^{-11}), \\
B(K_L \rightarrow \pi^0 \mu^\pm e^\mp) &\sim 1 \times 10^{-11} &< 7.6 \times 10^{-11}), \\
B(D^+ \rightarrow \pi^+ \mu^- e^+) &\sim 6 \times 10^{-13} &< 3.4 \times 10^{-5}), \\
B(D^0 \rightarrow \pi^0 \mu^- e^+) &\sim 1 \times 10^{-13} &< 8.6 \times 10^{-5}), \\
B(B^+ \rightarrow K^+ \mu^- \tau^+) &\sim 2 \times 10^{-6} &< 7.7 \times 10^{-5}), \\
B(B^0 \rightarrow K^0 \mu^- \tau^+) &\sim 2 \times 10^{-6} &(\text{no data}).
\end{aligned} \tag{4.17}$$

We also predict $B(K_L \rightarrow \pi^0 \nu_e \bar{\nu}_\mu) \simeq B(K_L \rightarrow \pi^0 \mu e)/2$. We show the predicted branching ratios $B(P \rightarrow P' e_i \bar{e}_j)$ versus $v_F \equiv v_\Psi$ in Fig.1. Therefore, if v_Ψ is a few TeV, observations of the lepton-flavor violating K - and B -decays with $\Delta N_F = 0$ will be within our reach.

We may also expect to observe the lightest family gauge boson A_3^3 if its mass is a few TeV. For simplicity, we neglect the up- and down-quark mixings, i.e. $(u_1, u_2, u_3) \simeq (u, c, t)$ and $(d_1, d_2, d_3) \simeq (d, s, b)$. The observation is practically the same as that for Z' boson (for a review, see Ref.[11]). Although in the conventional Z' model, Z' couples to the fermions of all flavors, while the A_3^3 boson couples only to $\tau^+ \tau^-$, $\nu_\tau \bar{\nu}_\tau$, $b\bar{b}$ and $t\bar{t}$, so that the branching ratios are given by

$$B(A_3^3 \rightarrow \tau^+ \tau^-) \sim 2B(A_3^3 \rightarrow \nu_\tau \bar{\nu}_\tau) \sim \frac{1}{3}B(A_3^3 \rightarrow b\bar{b}) \sim \frac{1}{3}B(A_3^3 \rightarrow t\bar{t}) \sim \frac{2}{15}. \tag{4.18}$$

We may expect to observe a peak of $\tau^+ \tau^-$ (but no peak in $e^+ e^-$ and $\mu^+ \mu^-$) in $pp \rightarrow ggX \rightarrow A_3^3 X \rightarrow \tau^+ \tau^- X$ at the LHC and $e^+ e^- \rightarrow Z^*/\gamma^* \rightarrow A_3^3 X \rightarrow \tau^+ \tau^- X$ at the ILC, although these cross sections of the A_3^3 productions are small compared with that of the Z' production. Similar discussion can be applied to hadronic jets instead of $\tau^+ \tau^-$.

Finally, we would like to comment on a constraint from the observed K^0 - \bar{K}^0 mixing. As we stated previously, contributions from exchanges of the U(3) family gauge bosons to the K^0 - \bar{K}^0

mixing depend on the magnitudes of the family mixing U_d in the down-quark sector. At present, we know the observed values of the CKM mixing $V_{CKM} = U_d^\dagger U_d$, while we do not know the value of U_d . Tentatively, let us assume that the CKM mixing is dominantly given by the down-quark mixing, i.e. $U_d \simeq V_{CKM}$. Then, the dominant contribution comes from the exchange of the family gauge boson A_2^2 : $(g_F^2/2M_{22}^2)(V_{us}^*V_{cs})^2 = (z_2^2/4v_\Psi^2)(V_{us}^*V_{cs})^2 = (v_\Psi^2)^{-1} \times 6.76 \times 10^{-4}$. In the present model, the CP violating effect in the dominant contribution is negligibly small. Since the standard model gives $\Delta m_K^{SM} \simeq (7/6 - 5/6)\Delta m_K^{exp}$ [12] ($\Delta m_K^{exp} = (3.483 \pm 0.006) \times 10^{-12}$ MeV [5]), we consider that a contribution from new physics [13] is $|\Delta m_K^{NewPhys}| < \Delta m_K^{exp}/6$. If we assume the vacuum-insertion approximation, we obtain a constraint $v_\Psi \gtrsim \mathcal{O}(10^5)$ GeV, which suggests that the lightest gauge boson mass should also satisfy $m(A_3^3) \gtrsim \mathcal{O}(10^5)$ GeV. This result contradicts our speculation $r \sim 10^{-1}$. If this speculation is confirmed in future observations, we must build a quark mass matrix model with $U_d \simeq \mathbf{1}$ in the diagonal basis of the charged lepton mass matrix M_e , especially with $(U_d)_{12} \simeq 0$. (This means that the down-quark mass matrix M_d takes a similar structure except for a unit matrix term, i.e. $M_d \simeq k_d M_e + m_0 \mathbf{1}$.)

5 Concluding remarks

In conclusion, we have proposed a family gauge boson model with inverted hierarchical masses. The model has been motivated to give a SUSY scenario of the Sumino's cancellation mechanism between the radiative correction due to the photon and that due to the family gauge bosons. As stated in the end of Sec.1, the present model has many characteristics compared with the Sumino model: (i) It is easy to build a model that is anomaly free, since the model takes the canonical assignments of the U(3) family. (ii) The dangerous $\Delta N_F = 2$ interactions do not appear in the limit of no quark mixing. (iii) The family gauge bosons with the inverted hierarchical masses offer a new view for the low energy phenomenology. (iv) Since our model is based on a SUSY scenario, the VEV relations are kept (up to the SUSY breaking effects) although in this paper we did not discuss the derivations of the relation (1.1) and so on.

If we take the mass relation (1.1) seriously and we want to apply the Sumino mechanism to a SUSY scenario, the present model will be a promising model as an alternative one of the Sumino model. Whether the gauge boson mass hierarchy is inverted or normal will be confirmed by observing the direction of the deviation from the e - μ universality in the pure leptonic tau decays. The present experimental result, $R_\tau = 1.0017 \pm 0.0016$, is in favor of the inverted mass hierarchy although the error is still large. Since we speculate that the lightest gauge boson mass is a few TeV, we expect the deviation $\Delta R_\tau = R_\tau - 1 \sim 10^{-4}$. A tau factory in the near future will confirm this deviation. In addition, some lepton flavor violating K - and B -decays, e.g. $K^+ \rightarrow \pi^+ \mu^+ e^-$ and $B^+ \rightarrow K^+ \tau^+ \mu^-$, will be within our reach. We also expect a direct observation of $\tau^+ \tau^- / b\bar{b} / t\bar{t}$ decay modes while no excesses in the $e^+ e^- / \mu^+ \mu^-$ modes in the LHC and the ILC.

Acknowledgments

The authors would like to thank Y. Sumino for valuable and helpful conversations, and also M. Tanaka and K. Tsumura for helpful discussions for lepton flavor violating and collider phenomenology. One of the authors (YK) is supported by JSPS (No. 21540266).

References

- [1] T. Maehara and T. Yanagida, Prog. Theor. Phys. **60** (1978) 822; F. Wilczek and A. Zee, Phys. Rev. Lett. **42** (1979) 421.
- [2] Y. Sumino, Phys. Lett. B **671** (2009) 477.
- [3] Y. Sumino, JHEP **0905** (2009) 075.
- [4] Y. Koide, Lett. Nuovo Cim. **34** (1982) 201; Phys. Lett. B **120** (1983) 161; Phys. Rev. D **28** (1983) 252.
- [5] K. Nakamura *et al.* [Particle Data Group], J. Phys. G **37** (2010) 075021.
- [6] H. Arason, *et al.*, Phys. Rev. D **46** (1992) 3945.
- [7] T. Appelquist, Y. Bai and M. Piai, Phys. Lett. B **637** (2006) 245; Phys. Rev. D **74** (2006) 076001.
- [8] Y. Koide, Mod. Phys. Lett. A **5** (1990) 2319.
- [9] Y. Koide, Phys. Rev. D **79** (2009) 033009.
- [10] W. Fetscher and H.-J. Gerber, in Ref.[5], p.521.
- [11] J. L. Hewett and T. G. Rizzo, Phys. Rept. **183** (1989) 193; A. Leike, Phys. Rept. **317** (1999) 143.
- [12] T. Inami and C. S. Lim, Prog. Theor. Phys. **65** (1981) 297 [Erratum-ibid. **65** (1981) 1772].
- [13] Y. Bai, UMI-3293287.